

Sigma Signal for Hybrid Baryon Decay

Leonard S. Kisslinger

Department of Physics, Carnegie Mellon University

Pittsburgh, PA 15213, USA and

Los Alamos National Laboratory, Los Alamos, NM 87545, USA

and

Zhenping Li

Physics Department Peking University, Beijing 100871, P. R. China

February 1, 2008

Abstract

We develop an ansatz of the sigma enhancement of the $I=0$, $L=0$ π - π scattering amplitude as arising from a low-energy glueball pole. Using this picture we estimate the $\pi^0\pi^0$ to π^0 branching ratio for the decays of the Roper resonance, which we previously found to be a hybrid in our QCD sum rule calculation.

PACS numbers: 12.39.Mk, 11.55.Hx, 14.20.Gk

In an investigation of scalar glueballs one of us has found[1] that a pure scalar glueball has a mass 300-600 MeV, while solutions for mixed scalar glueballs and mesons are at a much higher energy, consistent with the $f_0(1500)$ for the mainly-glueball solution and with the $f_0(1370)$ and $a_0(1450)$ for the mainly-meson solution. Recently there has been a great deal of experimental activity in glueball searches[2, 3]. One of the very interesting and we feel most important observations is the enhancement of the σ mode in the decays of the glueball candidates, such as the dominance of the $\sigma\sigma$ intermediate states in the decays of the glueball candidates $f_0(1500) \rightarrow 4\pi$ and $f_0(1710) \rightarrow 4\pi$ [3, 4]. The recent analysis by the BES collaboration[4] also suggest that the tensor glueball candidates $\xi(2230)$ also has a large 4π branching ratio that is dominated by the $\sigma f_2(1270)$ intermediate state.

By the " σ " we mean in the present paper the enhancement in the $\pi - \pi$ $I=0, L=0$ amplitude, for which a pole has been found in the analysis of Zou and Bugg[5]. This broad pole occurs near the light scalar glueball found in our work[1], and the work of others using QCD sum rule methods[6]. We propose that the physical origin of the σ enhancement is the light scalar gluonic mode, and that this could be a good signal for the study of gluonic structure of hadrons. This also provides an explanation of the absence of light, narrow glueball in lattice gauge calculations, since the strong coupling to the $\pi - \pi$ channel, which results in about a 700 MeV width must be included for a physical solution. Note the the mixed glueball-meson states, predicted to be at higher energies in the region of scalar mesons[1, 7], are quite far from this coupled-channel sigma and would not be involved.

In the present work we use this picture of a scalar glueball mode coupled to the σ channel for the decay of hybrid baryons. Recently we carried out a QCD sum rule calculation[8] to find the hybrid baryon with quantum numbers $J^P = 1/2^+, I = 1/2$, the same as the nucleon and the $P_{11}(1440)$, the "Roper". This consisted as an

improvement over earlier work using the QCD sum rule methods[9, 10]. Since that work we have considered currents of the form $c_1\eta_N + c_2\eta_H$, where η_N and η_H are the nucleon and hybrid currents, defined below, and have shown that the nucleon has a very small hybrid component[11], which indicates that using a pure hybrid current for the Roper is reasonable.

Because of convenience in treating renormalization we use a different current operator for the $J^P = 1/2^+, I = 1/2$ hybrid state than in Ref [8]. It is

$$\eta_H(x) = \epsilon^{abc} \left(u^a(x) C \gamma^\mu u^b(x) \right) i \sigma^{\alpha\beta} \gamma^\mu \gamma^5 G_{\alpha\beta}^d(x) \left(T^d d(x) \right)^c, \quad (1)$$

where $u(x)$ and $d(x)$ are operators of the u and d quarks, $G_{\mu\sigma}^d(x)$ is a gluon field strength, $a, b, c = 1, 2, 3$ are the color indices, C is the charge conjugation matrix, and $T^d = \lambda^d/2$ is the generator of the $SU(3)$ color group. For the calculation of the pion and sigma decays we use the external field two-point method[12], which has been used in many calculations of hadronic coupling constants, including the strong and weak pion-nucleon coupling constant[13]. Using a two-point function in an external J_Γ field to represent the coupling of the current operator to the nucleon and the Roper, with the current of Eq. 1 lead to two independent invariant functions;

$$\begin{aligned} \Pi^{st}(q) &= i \int e^{iqx} \langle 0 | T \left(\eta_H^s(x) \bar{\eta}_N^t(0) \right)_\Gamma | 0 \rangle \\ &= \Pi_1^\Gamma(q^2) \hat{q}^{st} + \Pi_2^\Gamma(q^2) \delta^{st}, \end{aligned} \quad (2)$$

where s, t are Dirac indices, $\hat{q}^{st} = (\gamma_\mu)^{st} q^\mu$, and η_N is the nucleon current. The details of the solution for the mass of the hybrid are given in Ref [8].

The coefficients of the operator product expansion (O.P.E.) are calculated to dimension 6. In addition to the leading perturbative contributions we include the two quark condensate terms, which are the largest nonperturbative terms. For the pion decay the current $J_\pi = i\bar{q}\gamma_5 q$ and the Π_2 correlator are used. After the Borel transform

we find the following sum rule

$$\begin{aligned}\Pi_2^\pi(M^2) &= G_\pi \frac{\lambda_N \lambda_H}{(M^N + M^H)} (M_H e^{-M_H^2/M^2} - M_N e^{-M_N^2/M^2}) (2\pi)^4 \\ &= \frac{11}{12} M^8 E^3 - \frac{4a^2}{9} M^2 E_0,\end{aligned}\quad (3)$$

where λ_N , λ_H are the QCD sum rule structure constants, and G_π is the pi-Roper-nucleon coupling. The susceptibility term is small and has been neglected. For the sigma decay we use the current J_σ , with the glue-sigma coupling, g_σ , given by

$$\langle G_{\alpha\beta}^a J_\sigma G_a^{\alpha\beta} \rangle = g_\sigma \langle G_{\alpha\beta}^a G_a^{\alpha\beta} \rangle. \quad (4)$$

The glue-sigma coupling constant can be extracted from the $\pi - \pi$ T-Matrix of Ref [5]. After subtracting the higher resonances they find a fit to the $\pi - \pi$ amplitude of a Breit-Wigner form:

$$T^{\pi\pi(\sigma)} = \frac{\sqrt{s} \Gamma_\sigma(s)/2}{s - M_\sigma^2 + i\sqrt{s} \Gamma_\sigma(s)/2} + T(background). \quad (5)$$

The analysis of Ref [5] finds the sigma pole at 370 - 356i. Using our ansatze we determine the sigma- gluonic coupling constant, g_σ from the width of this pole, giving

$$g_\sigma \simeq 700 MeV. \quad (6)$$

We obtain for the Π_2 correlator

$$\begin{aligned}\Pi_2(M^2) &= G_\sigma \frac{\lambda_N \lambda_H}{(M^H - M^N)} (e^{-M_N^2/M^2} - e^{-M_H^2/M^2}) (2\pi)^4 \\ &= g_\sigma \left(\frac{7}{30} M^8 E^3 - \frac{a^2}{3} M^2 E_0 \right).\end{aligned}\quad (7)$$

In Eq.(7) the parameter G_σ is the H-N- σ coupling constant which appears in the numerator of the double-pole term in the phenomenological dispersion relationship.

From G_σ and G_π the π to σ branching ratio can be determined. Using the standard values of the parameters, we find for the σ/π branching ratio

$$\frac{\Gamma_\sigma}{\Gamma_\pi} \simeq 10\% \cdot R_p, \quad (8)$$

where R_p is the ratio of phase space factors. This is consistent with the present experimental limits[14]. Indeed, the data suggest that the width of the Roper resonance $P_{11}(1440)$ decaying into σN final states is generally an order of magnitude larger than those of other resonances. This could be another signature of a hybrid Roper resonance in addition to its electromagnetic transition properties[15]. Thus, further experimental investigations of $P_{11}(1440) \rightarrow \pi^0 \pi^0 N$ [to select the $I=0$ channel and eliminate Δ and ρ backgrounds] over an energy range to map out the σ would be a very important channel to study the structure of the Roper resonance. In general, we suggest that the measurement of σ decays of baryons and mesons is a signal for the gluonic content of hadrons.

We would like to thank Mikkel Johnson for helpful discussions. This work was supported by National Science Foundation grants PHY-9319641 and INT-9514190, and by the Department of Energy.

References

- [1] L.S. Kisslinger, J. Gardner and C. Vanderstraeten, "Mixed Scalar Glueballs and Mesons", to be published in Phys. Lett. B (1997).
- [2] Crystal Barrel Collaboration, Phys. Lett. **B 355**, 425 (1995).
- [3] D. V. Bugg, *et al*, Phys. Lett. **353B**, 378 (1995).
- [4] Zhu Yucan, BES J/ Ψ group, to appear on The Proceedings of the 6th Conference On The Intersection of Particle and Nuclear Physics, Big Sky, Montana (1997).
- [5] B.S. Zou and D.V.Bugg, Phys. Rev. **D 50**, 591 (1994).

- [6] C.A. Dominguez and N. Paver, Zeit. Phys. **C 31**, 591 (1986); J. Bordes, V. Gimenez and J.A. Penarrocha, Phys. Lett **B 223**, 251 (1989); J. Liu and D. Liu, J. Phys. G:Nucl. Part. Phys. **19**, 373 (1993).
- [7] S. Narison, hep-ph/9612457.
- [8] L.S. Kisslinger and Z. Li, Phys. Rev. **D 51**, R5986 (1995).
- [9] A.P. Martynenko, Sov. J. Nucl. Phys. **54**, 488 (1991).
- [10] V.M. Braun, P. Gornicki, L. Mankiewicz and A. Schafer, Phys. Lett. **B 302**, 291 (1993).
- [11] L.S. Kisslinger, "Gluonic Hadrons", to be published in the Proceedings of the International Conference on Quark Lepton Nuclear Physics, Osaka, Japan (1997).
- [12] B.L. Ioffe and A.V. Smilga, Nucl. Phys. **B 232**, 109 (1984).
- [13] E.M. Henley, W-Y. P. Hwang and L.S. Kisslinger, Phys. Lett. **B 367**, 21 (1996).
- [14] Review of Particle Properties, Phys. Rev. **D 45** (1992).
- [15] Zhenping Li, V. Burkert and Zhujun Li, Phys. Rev. **D46**, 70 (1992); Zhenping Li, Phys. Rev. **D44**, 2841 (1991).